

# OMIC

OPHTHALMIC MUTUAL  
INSURANCE COMPANY  
(A Risk Retention Group)

655 Beach Street  
San Francisco CA 94109-1336

P.O. Box 880610  
San Francisco CA 94188-0610

Phone: (800) 562-6642, ext. 639  
Fax: (415) 771-7087  
Email: [omic@omic.com](mailto:omic@omic.com)  
Web site: [www.omic.com](http://www.omic.com)

OMIC believes that prudent refractive surgery underwriting requirements help protect both the company and its insureds. These requirements, based on sensible medical practice and sound risk management principles, were developed by practicing refractive surgeons to reduce the likelihood of claims and to aid in the defensibility of any resulting claims. OMIC is proud that its claims experience is significantly better than the industry average, and we believe this is due in large part to the company's underwriting requirements.

OMIC routinely reviews its refractive surgery requirements and modifies them when warranted. To date, nearly all revisions have served to expand, rather than reduce or restrict, coverage.

### APPLICABLE TO ALL PROCEDURES

- Prior to surgery, the surgeon must perform and document an independent evaluation of the patient's eligibility for surgery, including performing a slit lamp exam and reviewing topography, pachymetry, pupil size, and refractive stability, and discuss monovision option for presbyopic patients.
- Patient age: at least 18 years old, stable refraction without contact lenses (off-label for Summit patients <21) for RK/AK, PRK, LASIK and other accepted variations, and Intacs; over 40 years old, stable refraction without contact lenses for CK.
- Operating surgeon must conduct informed consent discussion.
- Surgeon must document in medical record that risks, benefits, alternatives, and complications were discussed.
- Each patient must be offered a copy of the consent form prior to the day of surgery.
- Surgeon or a designated ophthalmologist must perform the first postoperative visit.
- Advertisements (print, audio, video, Internet) must not be misleading or guarantee results.
- Surgeon must follow FDA and FTC advertising guidelines.

### LASIK/PRK/LASEK/INTRALASIK/CUSTOM-CAP AND OTHER ACCEPTED VARIATIONS

- Surgeon must be certified on laser to be used.
- Surgeon must be appropriately trained on the microkeratome (if applicable).
- Criteria for degree of myopia, hyperopia, and astigmatism must fall within FDA-approved guidelines. Off-label treatment of up to 6.0D astigmatism, -15.0D myopia, and +6.0D hyperopia permitted subject to special consent language.
- **First 5** cases must be performed **unilaterally** with minimum interval of one day. Bilateral simultaneous requires sufficient prior experience + **special OMIC consent form**. No off-label use permitted for bilateral simultaneous PRK.
- Retreatment permitted after documented refractive stability.

### RADIAL AND ASTIGMATIC KERATOTOMY

- Astigmatism: AK alone 1.0 D to 4D (demonstrate over-correction if <1.5 D), AK with RK up to 6D (otherwise demonstrate residual to patient).
- Myopia: -1D to -8D (demonstrate residual if more than 6D).
- Bilateral simultaneous not covered for primary surgery.
- Minimum of one week between primary procedure on first and second eye and between primary procedure and enhancement on same eye.

## INTRASTROMAL CORNEAL RING SEGMENTS ("INTACS")

- Astigmatism:  $\leq 1.0D$ ; Myopia:  $-1D$  to  $-3D$ .
- Bilateral simultaneous requires previous unilateral Intacs experience (5 cases), **special OMIC consent form**.
- Underwriting requirements apply only to treatment of myopia. Application and underwriting review not required if performance of Intacs procedures is limited to therapeutic purposes (e.g., keratoconus, post-LASIK ectasia).

## CONDUCTIVE KERATOPLASTY (CK)

- Astigmatism:  $\leq 0.75D$  ; Hyperopia:  $0.75D$  to  $3.0D$ .
- Presbyopia:  $\leq 0.75D$  astigmatism, emmetropia or mild hyperopia; spherical hyperopic treatment of  $1$  to  $2.25D$ ; successful trial or history of monovision; special requirements for treatment of dominant eye.
- Bilateral simultaneous requires **special OMIC consent form**.

## REFRACTIVE LENS EXCHANGE

- Myopia:  $-6 D$  to  $-15 D$ . Patients must be age  $40+$  and presbyopic.
- Hyperopia: Axial length  $20$  mm or longer. UCVA  $20/40$  or worse.  $+1 D$  to  $+15 D$  if age  $40+$  and presbyopic.  $+4 D$  to  $+15 D$  if under age  $40$ .
- Coverage not available for treatment of emmetropic patients (with or without presbyopia).
- Patients must undergo complete retinal exam and be advised of increased risk of retinal detachment.
- The procedure must be performed in a hospital or outpatient surgery center approved for cataract surgery. Full sterile technique must be followed.
- Minimum interval of one week between primary procedures.

## PHAKIC IMPLANTS FOR REFRACTIVE PURPOSES

- For FDA-approved implants, criteria for age, degree of myopia and astigmatism, and for anterior chamber depth must fall within FDA-approved guidelines.
- For cases performed during course of clinical trials, patients must fall within patient selection criteria established by trial protocol.
- The procedure must be performed in a hospital or outpatient surgery center approved for cataract surgery. Full sterile technique must be followed.
- Minimum interval of one week between primary procedures.

This abbreviated list is provided as a quick reference for OMIC insureds and is not a complete list of underwriting requirements or refractive procedures covered. Coverage for refractive surgery is not automatic and must be specifically applied for. Call OMIC's Underwriting Department at **(800) 562-6642**, send us an email at **[underwriting@omic.com](mailto:underwriting@omic.com)**, or access our web site at **[www.omic.com](http://www.omic.com)** for clarification or more detailed information.